



TITLE:

# Multi-polar orders in spin-1/2 frustrated magnets(Topological Aspects of Solid State Physics)

AUTHOR(S):

Momoi, Tsutomu

---

CITATION:

Momoi, Tsutomu. Multi-polar orders in spin-1/2 frustrated magnets(Topological Aspects of Solid State Physics). 物性研究 2009, 91(6): 674-674

ISSUE DATE:

2009-03-20

URL:

<http://hdl.handle.net/2433/142922>

RIGHT:

DAY 1: 13:30 – 14:10

## Topological Aspects of RNA Folding

Anthony Zee

KITP, UCSB

I will describe an approach to RNA folding using the topological features of large  $N$  matrix theory. After a brief review of the relevant biochemistry, I will describe the matrix theory and recent numerical work.

---

DAY 1: 14:10 – 14:50

## Multi-polar orders in spin-1/2 frustrated magnets

Tsutomu Momoi

RIKEN

We present a new scenario for the breakdown of ferromagnetic order in spin-1/2 frustrated magnets, which gives rise to “spin liquid” phases with hidden magnetic multi-polar orders. Dynamical effects lead to the formation of magnon bound states, which undergo Bose-Einstein condensation. This scenario is explored in some detail for Heisenberg models on the 2D square and triangular lattices, and also on the 1D chain.

On the square lattice, two-magnon bound states are most stable, giving rise to bond-centered spin nematic - i.e. quadrupolar - order but no long-range spin order in the ferromagnetic  $J_1$ - $J_2$  model. We present numerical evidence confirming the existence of a state with d-wave nematic correlations but no long-range spin order, lying between the saturated ferromagnetic and collinear antiferromagnetic phases [1]. In a model with ring exchange interactions on the triangular lattice, which is applicable to thin films of solid  $^3\text{He}$ , three-magnon bound states are most stably formed. Condensation of these three magnon bound states leads to a gapless spin liquid with hidden octupolar order in applied magnetic field [2].

In the 1D Heisenberg chain with competing ferromagnetic nearest-neighbor and antiferromagnetic next-nearest neighbor interactions, the number of magnons forming the bound states increases consecutively with approaching the ferromagnetic phase boundary, which leads to a cascade of phase transitions. Bose condensation of the bound  $n$  magnons below the saturation field leads to novel Tomonaga-Luttinger liquids with multi-polar correlations [3].

[1] N. Shannon, T. Momoi, and P. Sindzingre, Phys. Rev. Lett. **96**, 027213 (2006).

[2] T. Momoi, P. Sindzingre, and N. Shannon, Phys. Rev. Lett. **97**, 257204 (2006).

[3] L. Kecke, T. Momoi, and A. Furusaki, Phys. Rev. B **76**, 060407(R) (2007); T. Hikihara, L. Kecke, T. Momoi, and A. Furusaki, preprint.